

WINDOW EXPLOITATION STUDIES

- ++++ 8/7/77. Exp. to see if sensitivity to LSD is reduced (protected against) by pretreatment with MDMA-type drugs - vis., the ± of 7/4/77 at 60 µg on the day following 160 mg MDIP and 80 mg MDMA. At [3:35] into 200 mg MDE (post-symptomatic) take 60 µg LSD. =[0:00] [0:40 - 1:00] build up of sensory sparkle - mentally clear. [1:20] development into ++ of LSD with edge-loss of reality. This is as clear an LSD as I have ever in my life experienced. The temptation is to propel it deeper and more open with another 40 to 60 µg. No - I am too much on the rim of reality as it is - learn to in-and-out with this. [1:50] The world is dancing around me. Terrible temptation to expand the experiment. RESIST. [1:54] I am at a place, fully +++, where neither the MD series or LSD alone can get me. When LSD has to do the opening job, it pretty well tires you out - but with MDE to open things? is MDE needed - could MDMA work? Might this allow the detection of v. small amts. of LSD? Let's next try 120 MDMA and 60µg LSD together, or slightly spaced. I am at ++++ - I must resist the temptation to supplement and extend this state, but I don't want it to end. [2:40] perhaps the slight sign of dropping. at [3:00] am at ++, and resist the 40 µg additional LSD all ready! [4:00] at full happy + - and I am not yet sure if I am glad or sorry I resisted. That was probably the most surely tempting maneuver that I have ever been confronted with. At my present + I am complete ecstatic about my experiment. [7:00] some physical residues. The dancing, beautiful, in love with everything world of the [2:00 - 2:30] period was one of my rare ++++ peak experiences. Will I ever succeed in recapturing it? Let us try 120 mg MDMA and at about 1 hr, 60 µg LSD. Evening - excellent sleep. no AM res.
- + 8/10/77 MDMA -> LSD. 120 mg MDMA at [-2:00] q.v. 60 µg LSD at [0:00] [0:35] quiet window to about [1:20] to a + at most. [1:30] ho hum window - outside no better [2:00] dropping MDMA does not do it [2:40] virtually out - the two drugs do not see one-another
- +++ 8/14/77 MDE -> LSD 200 mg MDE at [-3:00] q.v. 60µg LSD=[0:00] notice at [0:30] - slow and quiet development to + at [0:45] [1:05] further from [1:00 to 2:00] very enjoyable develop. to a full +++ [2:30] spectacular world - I would like to maintain or extend this state. [3:00] still there [3:45] back to a + or ++ - next time lets try 200 mg MDE + 60 µg LSD together. [4:30] still big + and happy [5:30] still aware - no sleep problems.
- ++ 8/17/77 MDIP -> LSD 250 mg MDIP at [-2:30] q.v. 60 µg LSD=[0:00] [0:35] notice first edge effect [0:45] slow development [1:00] up to +, still devel. [1:15] going into what would be LSD alone. [2:00] straight LSD at ++ [2:30] drive to red-wood grove, back. edge-reality loss - if there is any enhancement of LSD - it is in the direction of reality loss, NOT window. [3:15] dropping [5:00] dropping, but am still +; [8:00] still trace residual. at [12:00] to bed - no sleep to [16:00], there only fretful. AM - fine, no residue

cont - 249