



4-thiomescaline. f. page 239.

+++ 40mg 12/8/79 5:38=[0:00] AP≡. 7 days clear. alert[:22] - developing [:30] - quite a bit of physical concern - I - malaise, not nausea, AP heavy tremor. Well developed at [1:30] to [2:00]. Complete thought mid-directions - some eroticism, but preference to talk, thought möbius reality changing. Shakes of DOB. [3:00] clock stare - [6:00] still +++ - scant urine search for fluids [7:00-9:00] AP some sleep, ATS little if any. [9:30] still +, wine to sleep. overall +++, with much convoluted thought process. Next day, somewhat depleted, selfish mood. Not a drug for fun, but intensely powerful. Not for group. Must be one on one - too much chance for psychological abreaction. Use care.

++ 30mg 2/1/80 (AG, MP ≡). Onset of alert, more at [:30 to 1:00] - very quiet build up to peak at [2:00 or 2:30]. Eyes closed fantasy involving body viewing, mobile structures rather than scenes. AG drops quickly, then recognizes differences in effects from expected. TG at [4:00] - sees cognition sobriety, but recognizes easy in talk (AG vs MP). Fantasy skill slipping at [6:00] - below + by [8:00]. Extremely chilled (internally) despite 72° house. Repeat in hot weather, at 30 or 40. This ++ at most.

25-40mg-active range
for thiomescaline paper

++(.5) 25mg 11/22/80 ATS, AP≡; NT, CT. 20mg 11:25AM=[0:00] - Alerts all [0:35 ±5] quiet buildup to [1:15-1:30] - clearly ++ or above - max [2:00-~4:00] quiet decline to [8:00-10:00] ATS - somewhat >+ - hodgepogy fantasy - some interesting - some ugly - disjointed thought processes - good, long sleep - no AM residue. NT accepted excellently - no concern - felt to be good experiment - some fantasy. CT [with] extreme tremors, at ~[1:00-1:30] and again ~[4:00] - floating feeling on walking. limber - good sleep. AP still ++ at [7:00] - very tired - good sleep. NT trouble sleeping. AM - all rested, no depletion. most ~+++ me, >+. All [with] profound anesthesia - perhaps light anorexia.