



stray information from here and there


cont: p 652.




ex. Wes Rusher 50mg stimulating, up all night, no MDA like effects. letter 5/26/81




ex. E. Presgraves 9, and 11 mg → +++ short lived




ex. E. Presgraves 3mg ± , 5mg - v.hygroscopic.



ex. Wes R. less active than the 5-OCH₃ (12/81)
4.5mg - at [:15]
+4.5 → Σ 8-10mg




ex. Zole Abrego N of 2, at 100mg, ±
N of 2, at 250mg, all >±, 8hrs
(1 [with] changes of cognitive function, later claimed "no effects").




ex. Dong Embree 20mg i.v. → effects in a couple of seconds. MDMA-like, relaxing, PSOP visuals

8 trys 35-40mg? orally - more relaxed, less demanding than 2C-B - compare closely [with] MDMA.




ex. Ruban Hubbell orally active at 500-750mg - visual - lasts 12 hrs.




280 S.Virginia, Reno
NS+NaoMe + reduction
see page 628


ex. Neal Aquiniga (orally?) active, less so than 2C-D, but yet more effective in opening up people.
his 2C-D 10mg, this 15mg - called MLO snort ~10mg. HCl rough, tartrate easier.
all [with] 3% Na[Hg] in MeOH? +HOAc?



ex. EH No effects orally at 30mg



ex. EH Active 8-10mg. EH [with] ++@10 - see page 625



ex. Neal Aquiniga 15mg orally; ≅ 2CD, taste ↑, smell ↓
- has made OEt - no assay
(+Wendy) see page 628 - see 652 for MD analog



R = n-Propyl 20mg ~(-)
R = ethyl 20mg < Psilocin } in my notes from the phone ascribed to DF (Franklin?) check with him